

PubMed Nucleotide

Protein

Genome

Structure

PopSet

Taxonomy

OMIM

Bo

Search PubMed



for



Limits

Preview/Index

History

Clipboard

Details

About Entrez



Abstract



Sort



Save



Clip Add



Order

Text Version

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

E-Utilities

PubMed Services

Journals Database

MeSH Browser

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

LinkOut

Cubby

Related Resources

Order Documents

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

Privacy Policy

☐ 1: Expert Opin Biol Ther 2002 Jan;2(1):3-24

Related Articles, Links



HSP90 as a new therapeutic target for cancer therapy: the story unfolds.

Workman P, Maloney A.

CRC Centre for Cancer Therapeutics, Institute of Cancer Research, Block E, 15 Cotswold Road, Sutton, Surrey SM2 5NG, UK. paulw@icr.ac.uk


Current anticancer drug development strategies involve identifying novel molecular targets which are crucial for tumourigenesis. The molecular chaperone heat shock protein (HSP) 90 is of interest as an anticancer drug target because of its importance in maintaining the conformation, stability and function of key oncogenic client proteins involved in signal transduction pathways leading to proliferation, cell cycle progression and apoptosis, as well as other features of the malignant phenotype such as invasion, angiogenesis and metastasis. The natural product HSP90 inhibitors geldanamycin and radicicol exert their antitumour effect by inhibiting the intrinsic ATPase activity of HSP90, resulting in degradation of HSP90 client proteins via the ubiquitin proteasome pathway. Anticancer selectivity may derive from the simultaneous combinatorial effects of HSP90 inhibitors on multiple cancer targets and pathways. 17-allylamino, 17-demethoxygeldanamycin (17AAG), a geldanamycin derivative, showed good activity and cancer selectivity in preclinical models and has now progressed to Phase I clinical trial in cancer patients with encouraging initial results. Phase II trials including combination studies with cytotoxic agents are now being planned and these should allow the therapeutic activity of 17AAG to be determined. Second generation HSP90 inhibitors may be designed to overcome some of the drawbacks of 17AAG, including limited oral bioavailability and solubility. They could also be engineered to target specific functions of HSP90, which may not only provide greater molecular selectivity and clinical benefit but may also increase understanding of the complex functions of this molecular chaperone. HSP90 inhibitors provide proof of concept for drugs directed at HSP90 and protein folding and this principle may be applicable to other medical conditions involving protein aggregation and stability.

PMID: 11772336 [PubMed - in process]

Display	Abstract	<input type="checkbox"/>	Sort	<input type="checkbox"/>	Save	Text	Clip Add	Order
---------	----------	--------------------------	------	--------------------------	------	------	----------	-------

[Write to the Help Desk](#)
[NCBI](#) | [NLM](#) | [NIH](#)
[Department of Health & Human Services](#)
[Freedom of Information Act](#) | [Disclaimer](#)

i686-pc-linux-gnu Oct 31 2002 15:09:13



PubMed Nucleotide Protein Genome Structure PopSet Taxonomy OMIM Bo

Search for



Limits Preview/Index History Clipboard Details

About Entrez



Text Version

Entrez PubMed
Overview
Help | FAQ
Tutorial
New/Noteworthy
E-Utilities

PubMed Services
Journals Database
MeSH Browser
Single Citation Matcher
Batch Citation Matcher
Clinical Queries
LinkOut
Cubby

Related Resources
Order Documents
NLM Gateway
TOXNET
Consumer Health
Clinical Alerts
ClinicalTrials.gov
PubMed Central

Privacy Policy

☐ 1: Cancer Res 2001 Apr 1;61(7):2945-52

Related Articles, Links

FREE full text article at
cancerres.aacrjournals.org


Inhibition of heat shock protein 90 function by ansamycins causes the morphological and functional differentiation of breast cancer cells.

Munster PN, Srethapakdi M, Moasser MM, Rosen N.

Department of Medicine, Memorial Sloan-Kettering Cancer Center, New York, New York 10021, USA.

17-(Allylamino)-17-demethoxygeldanamycin (17-AAG) is an ansamycin antibiotic that binds to a conserved pocket in Hsp90 and induces the degradation of proteins that require this chaperone for conformational maturation. 17-AAG causes a retinoblastoma (RB)-dependent G1 block in cancer cells and is now in clinical trial. In breast cancer cells, G1 block is accompanied by differentiation and followed by apoptosis. The differentiation is characterized by specific changes in morphology and induction of milk fat proteins and lipid droplets. In cells lacking RB, neither G1 arrest nor differentiation occurs; instead, they undergo apoptosis in mitosis. Introduction of RB into these cells restores the differentiation response to 17-AAG. Inhibitors of the ras, mitogen-activated protein kinase, and phosphatidylinositol 3-kinase pathways cause accumulation of milk fat proteins and induction of lipid droplets when associated with G1 arrest but do not cause morphological changes. Thus, regulation of Hsp90 function by 17-AAG in breast cancer cells induces RB-dependent morphological and functional mammary differentiation. G1 arrest is sufficient for some but not all aspects of the phenotype. Induction of differentiation may be responsible for some of the antitumor effects of this drug.

PMID: 11306472 [PubMed - indexed for MEDLINE]



Write to the Help Desk
NCBI | NLM | NIH
Department of Health & Human Services

Freedom of Information Act | Disclaimer

i686-pc-linux-gnu Oct 31 2002 15:09:13